

# 5. FEEDING ACTIVITY IN THE CAPTIVITY OF THE WESTERN GHATS SQUIRREL *FUNAMBULUS TRISTRIATUS* WATERHOUSE<sup>1</sup>

(With a text-figure)

## INTRODUCTION

The Western Ghats squirrel (*Funambulus tristriatus* Waterhouse) is an important rodent pest of cacao (*Theobroma cacao* L.) in south India (Bhat *et al.* 1981). But no effective

succeed the schedule of baiting must coincide with the peaks of feeding of the target animal. In this report the observations made by us on the rhythm of feeding of the Western Ghats squirrel are discussed.




Fig. 1. Feeding pattern of *Funambulus tristriatus*.

method has been evolved so far for its control. Trapping by offering baits is one of the methods usually tried for controlling this squirrel. If this type of control of rodent pests is to

## MATERIAL AND METHODS

The study was undertaken in the laboratory of the Central Plantation Crops Research Institute, Regional Station, Vittal, Dakshina Kan-nada district of Karnataka under natural day-light in November 1978. Six adult Western Ghats squirrels (3 ♂ ♂ and 3 ♀ ♀) of recorded

<sup>1</sup> Part of the thesis submitted to the University of Calicut by the first author for the award of Ph.D. degree 1983.

body weights were lodged individually in cells of  $50 \times 30 \times 30$  cm. These squirrels were allowed to acclimatise themselves to the conditions in the cage for five days prior to the experiment. On the sixth day each squirrel was provided with weighed quantity of paddy grains, previously air dried to a constant weight. The experiment was continued for 10 days. Each day the consumption of paddy was recorded at two-hourly intervals between 0400 and 2000 h. The weighing was done to the nearest 0.5 g using a common counter balance. Absolute consumption values were transformed to g/100 g body weight of the animal. The average consumption during each period was computed and analysed statistically.

#### RESULTS AND DISCUSSION

The data (see Fig. 1) revealed the presence of distinct bimodal feeding pattern in this squirrel. Feeding was first observed around 0600 h, increasing rapidly thereafter. The over-

all mean consumption per animal per two-hour-period was  $0.54 \pm 0.02$  g. The morning peak in feeding ( $1.44 \pm 0.09$  g) was observed around 0800 h. After 0800 h the feeding activity lessened gradually reaching a minimum ( $0.18 \pm 0.08$  g) at about 1400 h. Feeding activity increased again gradually after 1400 h reaching a peak around 1800 h. Thereafter feeding slowed down and ceased at about 2000 h. Feeding was never observed at night. It is evident from this study that baits for these squirrels are best set up in the early hours of the day.

#### ACKNOWLEDGEMENTS

We are grateful to Dr N. M. Nayar, former Director, CPCRI, Kasaragod, for laboratory facilities at CPCRI (Regional Station) Vittal and Shri B. P. Nair CPCRI (Regional Station, Vittal) for statistical analysis. One of us (SKB) is thankful to the Council of Scientific and Industrial Research, New Delhi, for a fellowship.

DEPARTMENT OF ZOOLOGY,  
UNIVERSITY OF CALICUT,  
KERALA-673 635,  
May 16, 1984.

S. KESHAHA BHAT<sup>1</sup>  
D. N. MATHEW

#### REFERENCE

BHAT, S. K., NAIR, C. P. R. & MATHEW, D. N.  
(1981): Mammalian pests of cocoa in south India.  
*Trop. Pest Mgmt.* 27:297-307.

<sup>1</sup> *Present Address:* Central Plantation Crops  
Research Institute, Reg. Stan., Kayangulam, Kerala-  
690 533.

#### 6. REPORT ON THE OCCURRENCE OF THE FAWN-COLOURED MOUSE, *MUS CERVICOLOR CERVICOLOR* HODGSON, 1845 [RODENTIA: MURIDAE] IN THE ANDAMAN AND NICOBAR ISLANDS, INDIA

A small collection of rodents obtained by of India, from the Andaman and Nicobar  
Shri P. K. Das, Zoologist, Zoological Survey Islands included a grey-bellied mouse with